

On Completeness of Eigenfunctions of the Spectral Problem

A.Q. Akhmedov, I.Q. Feyzullayev

Abstract. In this paper is studied the spectral problem for a discontinuous second order differential operator with a spectral parameter in conjugation conditions, that arises by solving the problem on vibrations of a loaded string with the free ends. Asymptotic formulas for the eigenvalues and eigenfunctions of the spectral problem are obtained, a theorem on the completeness and minimality of system of eigenvectors of the linearize operator in spaces $L_p \oplus C$ is proved, as well as a theorem on the completeness and minimality in L_p of the system of eigenfunctions of the spectral problem after excluding an arbitrary eigenfunction from it.

Key Words and Phrases: eigenvalues, eigenfunctions, asymptotic formula, biorthogonal system, completeness, minimality.

2010 Mathematics Subject Classifications: Primary 68N19, 68P10, 68T20

1. Introduction

Consider a spectral problem

$$y''(x) + \lambda y(x) = 0 \quad , x \in \left(0, \frac{1}{3}\right) \cup \left(\frac{1}{3}, 1\right), \quad (1)$$

$$\left. \begin{aligned} y'(0) = y'(1) = 0, \\ y\left(\frac{1}{3} - 0\right) = y\left(\frac{1}{3} + 0\right), \\ y'\left(\frac{1}{3} - 0\right) - y'\left(\frac{1}{3} + 0\right) = \lambda m y\left(\frac{1}{3}\right), \end{aligned} \right\} \quad (2)$$

which arises by solving the problem on vibrations of a loaded string with the free ends [1-3]. The spectral problem corresponding to the problem of oscillation of a loaded string with intersected ends was studied in [4;5] in the case when the load is fixed in the middle of the string, and in [6-8] in the case when the load is fixed at the point $x = \frac{1}{3}$. Similar questions for spectral problems corresponding to the case when the load is fixed at one or both ends of the string, were studied by other methods in [9-12].

2. The asymptotic of eigenvalues and eigenfunctions

Let $\lambda = \rho^2$ and introduce the following notation for boundary forms (2).

$$U_\nu(y) = U_{\nu 1}(y) + U_{\nu 2}(y), \nu = \overline{1, 4}, \quad (3)$$

where

$$U_{11}(y) = y'(0), U_{12}(y) \equiv 0,$$

$$U_{21}(y) \equiv 0, U_{22}(y) = y'(1),$$

$$U_{31}(y) = y\left(\frac{1}{3} - 0\right), U_{32}(y) = -y\left(\frac{1}{3} + 0\right),$$

$$U_{41}(y) = y'\left(\frac{1}{3} - 0\right), U_{42}(y) = -y'\left(\frac{1}{3} + 0\right) - \rho^2 m y\left(\frac{1}{3} + 0\right).$$

Firstly, let us prove the following theorem.

Theorem 1. *Spectral problem (1), (2) has two series of simple eigenvalues: $\lambda_{1,n} = (\rho_{1,n})^2$, $n = 1, 2, \dots$, è $\lambda_{2,n} = (\rho_{2,n})^2$, $n = 0, 1, 2, \dots$, where*

$$\left. \begin{aligned} \rho_{1,n} &= 3\pi n + \frac{3\pi}{2} + O\left(\frac{1}{n}\right) \\ \rho_{2,n} &= \frac{3\pi n}{2} + \frac{3\pi}{2} + O\left(\frac{1}{n}\right). \end{aligned} \right\} \quad (4)$$

The eigenfunctions $u(x)$, $n=0, 1, 2, \dots$, prescribed by formula

$$u_{i,n}(x) = \begin{cases} \cos \frac{2\rho_{i,n}}{3} \cos \rho_{i,n} x, & x \in \left[0, \frac{1}{3}\right], \\ \cos \frac{\rho_{i,n}}{3} \cos \rho_{i,n} (1-x), & x \in \left[\frac{1}{3}, 1\right], \end{cases} \quad i = 1, 2; n \in Z^+ \quad (5)$$

correspond to them.

Proof. Take $y_{11}(x) = \sin \rho x$, $y_{12}(x) = \cos \rho x$ for $x \in \left[0, \frac{1}{3}\right]$ and $y_{21}(x) = \sin \rho \left(x - \frac{1}{3}\right)$, $y_{22}(x) = \cos \rho \left(x - \frac{1}{3}\right)$ for $x \in \left[\frac{1}{3}, 1\right]$ as linear - independent solutions of the equation (1). Eigenfunctions of the problem (1), (2) will be sought in the form

$$\begin{aligned} y(x, \rho) &= \begin{cases} c_{11}y_{11}(x) + c_{12}y_{12}(x), & x \in \left[0, \frac{1}{3}\right], \\ c_{21}y_{21}(x) + c_{22}y_{22}(x), & x \in \left[\frac{1}{3}, 1\right], \end{cases} = \\ &= \begin{cases} c_{11}\sin \rho x + c_{12}\cos \rho x, & x \in \left[0, \frac{1}{3}\right], \\ c_{21}\sin \rho \left(x - \frac{1}{3}\right) + c_{22}\cos \rho \left(x - \frac{1}{3}\right), & x \in \left[\frac{1}{3}, 1\right]. \end{cases} \end{aligned} \quad (6)$$

Let us require that the function $y(x, \rho)$ satisfies the boundary conditions (2). Then for definition of numbers $c_{j,k}$ we obtain the system of the linear homogeneous equations

$$\begin{cases} C_{11}U_{\nu 1}(y_{11}) + C_{12}U_{\nu 1}(y_{12}) + C_{21}U_{\nu 2}(y_{21}) + C_{22}U_{\nu 2}(y_{22}) = 0, \\ \nu = \overline{1, 4}, \text{ (or briefly } \sum_{j,k=1}^2 C_{kj}U_{\nu j}(y_{jk}) = 0), \end{cases} \quad (7)$$

whose determinant is

$$\Delta(\rho) = \det \|U_{\nu j}(y_{jk})\|, j, k = 1, 2, \nu = \overline{1, 4}$$

Taking into consideration (3), for values of forms $U_{\nu j}(y_{jk})$ we have

$$\begin{cases} U_{11}(y_{11}) = \rho, U_{11}(y_{12}) = 0, U_{12}(y_{21}) = 0, U_{12}(y_{22}) = 0, \\ U_{21}(y_{11}) = 0, U_{21}(y_{12}) = 0, U_{22}(y_{21}) = \rho \cos \frac{2\rho}{3}, U_{22}(y_{22}) = -\rho \sin \frac{2\rho}{3}, \\ U_{31}(y_{11}) = \sin \frac{\rho}{3}, U_{31}(y_{12}) = \cos \frac{\rho}{3}, U_{32}(y_{21}) = 0, U_{32}(y_{22}) = -1, \\ U_{41}(y_{11}) = \rho \cos \frac{\rho}{3}, U_{41}(y_{12}) = -\rho \sin \frac{\rho}{3}, U_{42}(y_{21}) = -\rho, U_{42}(y_{22}) = -\rho^2 m \end{cases} \quad (8)$$

Opening determinant $\Delta(\rho)$, with the account (8) we obtain

$$\Delta(\rho) = \rho^3 \left(\sin \rho + \rho m \cos \frac{\rho}{3} \cos \frac{2\rho}{3} \right). \quad (9)$$

From the formula (9) it is obvious, that function $\Delta(\rho)$ has two series of zeros $\rho_{1,n}$ and $\rho_{2,n}$, which are asymptotically close to the zeros of the functions $\cos \frac{\rho}{3}$ and $\cos \frac{2\rho}{3}$, respectively.

Reasoning as in [13, p.20], we obtain that for $\rho_{1,n}$ and $\rho_{2,n}$ the following asymptotic formula are true

$$\rho_{1,n} = 3\pi n + \frac{3\pi}{2} + O\left(\frac{1}{n}\right), \rho_{2,n} = \frac{3\pi n}{2} + \frac{3\pi}{2} + O\left(\frac{1}{n}\right)$$

Taking into account (8), from (7) we obtain

$$c_{11} = 0,$$

$$c_{21} \cos \frac{2\rho}{3} - c_{22} \sin \frac{2\rho}{3} = 0,$$

$$c_{22} = c_{12} \cos \frac{\rho}{3}.$$

It is clear that in these relation $\cos \frac{2\rho}{3} \neq 0$ and choosing $c_{12} = \cos \frac{2\rho}{3}$, we get $c_{22} = \cos \frac{2\rho}{3} \cos \frac{\rho}{3}$, $c_{21} = \cos \frac{\rho}{3} \sin \frac{2\rho}{3}$. Taking into account the obtained values of the coefficients in (6), and substituting $\rho = \rho_{1,n}$ and $\rho = \rho_{2,n}$ there, respectively, we get that the eigenfunction $u_{i,n}(x)$, $i = 1, 2; n \in \mathbb{Z}^+$, corresponding to eigenvalue $\lambda_{i,n} = (\rho_{i,n})^2$ $i = 1, 2; n \in \mathbb{Z}^+$ in the following form

$$u_{i,n}(x) = \begin{cases} \cos \frac{2\rho_{i,n}}{3} \cos \rho_{i,n} x, x \in \left[0, \frac{1}{3}\right], \\ \cos \frac{\rho_{i,n}}{3} \cos \rho_{i,n} (1-x), x \in \left[\frac{1}{3}, 1\right], \end{cases} \quad i = 1, 2; n \in \mathbb{Z}^+$$

Theorem is proved.

3. Construction of the Green's function and the resolvents of the linearized operator

Now let's pass to construction of the Green's function of problem (1), (2). It is defined as a kernel of integral representation for solution of the corresponding non-homogeneous problem

$$y''(x) + \rho^2 y(x) = f(x), \quad (10)$$

satisfying boundary conditions (2). The solution of problem (10), (2) will be sought in the form

$$y(x) = \begin{cases} y_1(x), & \text{for } x \in [0, \frac{1}{3}] , \\ y_2(x), & \text{for } x \in [\frac{1}{3}, 1] , \end{cases} \quad (11)$$

where

$$\begin{cases} y_1(x) = c_{11}y_{11}(x) + c_{12}y_{12}(x) + \int_0^{\frac{1}{3}} g(x, \xi, \rho) f(\xi) d\xi, & x \in [0, \frac{1}{3}] , \\ y_2(x) = c_{21}y_{21}(x) + c_{22}y_{22}(x) + \int_{\frac{1}{3}}^1 g(x, \xi, \rho) f(\xi) d\xi, & x \in [\frac{1}{3}, 1] . \end{cases} \quad (12)$$

$$y_{11}(x) = e^{i\rho x}, y_{12}(x) = e^{-i\rho x}, \quad x \in [0, \frac{1}{3}] ;$$

$$y_{21}(x) = e^{i\rho(x-\frac{1}{3})}, y_{22}(x) = e^{-i\rho(x-\frac{1}{3})}, \quad x \in [\frac{1}{3}, 1]$$

$$g_1(x, \xi, \rho) = \begin{cases} -\frac{1}{2i\rho} (e^{i\rho(x-\xi)} - e^{-i\rho(x-\xi)}), & 0 \leq x < \xi \leq \frac{1}{3}, \\ \frac{1}{2i\rho} (e^{i\rho(x-\xi)} - e^{-i\rho(x-\xi)}), & 0 \leq x \leq \frac{1}{3}, \end{cases} \quad (13)$$

$$g_2(x, \xi, \rho) = \begin{cases} -\frac{1}{2i\rho} (e^{i\rho(x-\xi)} - e^{-i\rho(x-\xi)}), & \frac{1}{3} \leq x < \xi \leq 1, \\ \frac{1}{2i\rho} (e^{i\rho(x-\xi)} - e^{-i\rho(x-\xi)}), & \frac{1}{3} \leq \xi < x \leq 1. \end{cases} \quad (13')$$

Let us require that the function (11) satisfies the boundary conditions (2). Then for definition of numbers $C_{j,k}$ we obtain the system of algebraic equations

$$U_\nu(y) = \sum_{j,k=1}^2 C_{j,k} U_{\nu j}(y_{jk}) + \int_0^{\frac{1}{3}} U_{\nu 1}(g) f(\xi) d\xi + \int_{\frac{1}{3}}^1 U_{\nu 2}(g) f(\xi) d\xi = 0, \nu = \overline{1, 4}. \quad (14)$$

Let's define numbers $C_{j,k}$ from (14) and substituting their values in (12), for solving the problem (10), (12) we obtain the formula

$$\begin{aligned} y_1(x) &= \int_0^{\frac{1}{3}} G_{11}(x, \xi, \rho) f(\xi) d\xi + \int_{\frac{1}{3}}^1 G_{12}(x, \xi, \rho) f(\xi) d\xi, \quad x \in [0, \frac{1}{3}], \\ y_2(x) &= \int_0^{\frac{1}{3}} G_{21}(x, \xi, \rho) f(\xi) d\xi + \int_{\frac{1}{3}}^1 G_{22}(x, \xi, \rho) f(\xi) d\xi, \quad x \in [\frac{1}{3}, 1], \end{aligned} \quad (14')$$

where

$$G_{11}(x, \xi, \rho) = \frac{1}{\Delta(\rho)} \begin{vmatrix} g & y_{11} & y_{12} & 0 & 0 \\ U_{\nu 1}(g) & U_{\nu 1}(y_{11}) & U_{\nu 1}(y_{12}) & U_{\nu 2}(y_{21}) & U_{\nu 2}(y_{22}) \\ \dots & \dots & \dots & \dots & \dots \\ \nu=\overline{1,4} \end{vmatrix}$$

$$G_{12}(x, \xi, \rho) = \frac{1}{\Delta(\rho)} \begin{vmatrix} 0 & y_{11} & y_{12} & 0 & 0 \\ U_{\nu 2}(g) & U_{\nu 1}(y_{11}) & U_{\nu 1}(y_{12}) & U_{\nu 2}(y_{21}) & U_{\nu 2}(y_{22}) \\ \dots & \dots & \dots & \dots & \dots \\ \nu=\overline{1,4} \end{vmatrix}$$

$$G_{21}(x, \xi, \rho) = \frac{1}{\Delta(\rho)} \begin{vmatrix} 0 & 0 & 0 & y_{21} & y_{22} \\ U_{\nu 1}(g) & U_{\nu 1}(y_{11}) & U_{\nu 1}(y_{12}) & U_{\nu 2}(y_{21}) & U_{\nu 2}(y_{22}) \\ \dots & \dots & \dots & \dots & \dots \\ \nu=\overline{1,4} \end{vmatrix}$$

$$G_{22}(x, \xi, \rho) = \frac{1}{\Delta(\rho)} \begin{vmatrix} g & 0 & 0 & y_{21} & y_{22} \\ U_{\nu 2}(g) & U_{\nu 1}(y_{11}) & U_{\nu 1}(y_{12}) & U_{\nu 2}(y_{21}) & U_{\nu 2}(y_{22}) \\ \dots & \dots & \dots & \dots & \dots \\ \nu=\overline{1,4} \end{vmatrix}$$

and $\Delta(\rho)$ is a determinant from (7).

$$U_{11}(g) = -\frac{i\rho}{2i\rho} \left(e^{i\rho\xi} + e^{-i\rho\xi} \right), U_{12}(g) = 0$$

$$U_{21}(g) = 0, U_{22}(g) = \frac{i\rho}{2i\rho} \left(e^{i\rho(1-\xi)} + e^{-i\rho(1-\xi)} \right)$$

$$U_{31}(g) = \frac{1}{2i\rho} \left(e^{i\rho(\frac{1}{3}-\xi)} - e^{-i\rho(\frac{1}{3}-\xi)} \right), U_{32}(g) = \frac{1}{2i\rho} \left(e^{i\rho(\frac{1}{3}-\xi)} - e^{-i\rho(\frac{1}{3}-\xi)} \right) \quad (15)$$

$$U_{41}(g) = \frac{i\rho}{2i\rho} \left(e^{i\rho(\frac{1}{3}-\xi)} + e^{-i\rho(\frac{1}{3}-\xi)} \right), U_{42}(g) = \frac{i\rho}{2i\rho} \left(e^{i\rho(\frac{1}{3}-\xi)} + e^{-i\rho(\frac{1}{3}-\xi)} \right) +$$

$$+ \frac{\rho^2 m}{2i\rho} \left(e^{i\rho(\frac{1}{3}-\xi)} - e^{-i\rho(\frac{1}{3}-\xi)} \right)$$

$$U_{11}(y_{11}) = i\rho, U_{11}(y_{12}) = -i\rho, U_{12}(y_{21}) = 0, U_{12}(y_{22}) = 0,$$

$$U_{21}(y_{11}) = 0, U_{21}(y_{12}) = 0, U_{22}(y_{21}) = i\rho e^{\frac{2i\rho}{3}}, U_{22}(y_{22}) = -i\rho e^{-\frac{2i\rho}{3}},$$

$$U_{31}(y_{11}) = e^{\frac{i\rho}{3}}, U_{31}(y_{12}) = e^{-\frac{i\rho}{3}}, U_{32}(y_{21}) = -1, U_{32}(y_{22}) = -1,$$

$$U_{41}(y_{11}) = i\rho e^{\frac{i\rho}{3}}, U_{41}(y_{12}) = -i\rho e^{-\frac{i\rho}{3}}, U_{42}(y_{21}) = -i\rho - \rho^2 m, U_{42}(y_{22}) = i\rho - \rho^2 m.$$

Let's substitute (15), (8) and (9) in determinants of formula for $G_{kj}(x, \xi, \rho)$. Transforming the received determinants similar to [14, p. 95], and then opening them, we obtain the formula for the Green's function components. We'll formulate it as a lemma.

Lemma 2. *For the Green's function components $G_{kj}(x, \xi, \rho)$ of the problem (1), (2) the following expressions are true:*

$$\begin{aligned} G_{11}(x, \rho, \xi) = & \pm \rho^2 e^{i\rho(x-\xi)} - \frac{\rho^2}{\Delta(\rho)} \left(2e^{-i\rho} - i\rho m e^{-i\rho} = i\rho m e^{-\frac{i\rho}{3}} \right) e^{i\rho x} \cdot e^{i\rho \xi} - \\ & - \frac{\rho^2}{\Delta(\rho)} \left(2e^{i\rho} + i\rho m e^{-\frac{i\rho}{3}} + i\rho m e^{i\rho} \right) e^{i\rho x} \cdot e^{-i\rho \xi} - \end{aligned} \quad (16)$$

$$\begin{aligned} & - \frac{\rho^2}{\Delta(\rho)} \left(2e^{i\rho} + 2i\rho m e^{-\frac{i\rho}{3}} + i\rho m e^{i\rho} \right) e^{-i\rho x} \cdot e^{-i\rho \xi} - \\ & - \frac{\rho^2}{\Delta(\rho)} \left(2e^{i\rho} + i\rho m e^{i\rho} \right) e^{-i\rho x} \cdot e^{-i\rho \xi}, x \in \left[0, \frac{1}{3} \right], \xi \in \left[0, \frac{1}{3} \right], \end{aligned}$$

$$G_{12}(x, \rho, \xi) = 2e^{-\frac{2i\rho}{3}} e^{-i\rho x} e^{-i\rho(\xi-1)} \left(e^{\frac{2i\rho}{3}} - e^{i\rho(\xi-\frac{1}{3})} e^{i\rho(\xi-1)} \right) (e^{2i\rho x} + 1) \quad (17)$$

$$G_{21}(x, \rho, \xi) = 2e^{-i\rho} e^{-i\rho(\xi-\frac{1}{3})} e^{-i\rho(x-\frac{1}{3})} \left(e^{\frac{i\rho}{3}} e^{i\rho(\xi-\frac{1}{3})} e^{i\rho x} + 1 \right) \left(e^{\frac{4i\rho}{3}} + e^{2i\rho(x-\frac{1}{3})} \right) \quad (18)$$

$$\begin{aligned} G_{22}(x, \rho, \xi) = & \pm \rho^2 e^{i\rho(x-\xi)} + \frac{\rho^2}{\Delta(\rho)} \left((1 - i\rho m) e^{\frac{2i\rho}{3}} - (1 + i\rho m) e^{\frac{4i\rho}{3}} + e^{\frac{-2i\rho}{3}} \right) e^{i\rho(x-\frac{1}{3})} e^{-i\rho \xi} + \\ & + \frac{\rho^2}{\Delta(\rho)} \left(e^{-\frac{4i\rho}{3}} - (1 + i\rho m) e^{\frac{-2i\rho}{3}} - e^{\frac{4i\rho}{3}} \right) e^{i\rho(x-\frac{1}{3})} e^{i\rho \xi} - \end{aligned} \quad (19)$$

$$\begin{aligned} & - \frac{\rho^2}{\Delta(\rho)} \left(-(1 + i\rho m) e^{\frac{2i\rho}{3}} - (-3 + i\rho m) e^{\frac{4i\rho}{3}} \right) e^{-i\rho(x-\frac{1}{3})} e^{-i\rho \xi} - \\ & - \frac{\rho^2}{\Delta(\rho)} \left((1 + i\rho m) - (1 + i\rho m) e^{\frac{2i\rho}{3}} \right) e^{-i\rho(x-\frac{1}{3})} e^{i\rho \xi}, x \in \left[0, \frac{1}{3} \right], \rho \in \left[0, \frac{1}{3} \right]. \end{aligned}$$

Let us now proceed to the construction of linearizing operator. By $W_p^k(0, \frac{1}{3}) \oplus W_p^k(\frac{1}{3}, 1)$ we denote a space functions whose contractions on segments $[0, \frac{1}{3}]$ and $[\frac{1}{3}, 1]$ belong correspondingly to Sobolev spaces $W_p^k(0, \frac{1}{3})$ and $W_p^k(\frac{1}{3}, 1)$. Let's define the operator \mathcal{L} in $L_p(0, 1) \oplus C$ as follows:

$$D(\mathcal{L}) = \left\{ \hat{u} \in L_p(0, 1) \oplus C: \hat{u} = \left(u, mu \left(\frac{1}{3} \right) \right), u \in W_p^2 \left(0, \frac{1}{3} \right) \cup \left(\frac{1}{3}, 1 \right), \right. \\ \left. u'(0) = u'(1) = 0, u \left(-\frac{1}{3} \right) = u \left(\frac{1}{3} \right), \right\} \quad (20)$$

and for $\widehat{u} \in D(\mathcal{L})$

$$\mathcal{L}\widehat{u} = (-u''; u' \left(\frac{1}{3} - 0 \right) - u' \left(\frac{1}{3} + 0 \right)), \widehat{u} \in D(\mathcal{L}). \quad (21)$$

Lemma 3. *Operator defined by the formula (20), (21) is a linear closed operator with dense definitional domain in $L_p(0, 1) \oplus C$. Eigenvalues of the operator \mathcal{L} and problem (1), (2) coincide, and $\{\widehat{u}_k\}_{k \in N_0}$ are eigenvectors of the operator \mathcal{L} , where $N_0 = N \cup \{0\}$,*

$$\widehat{u}_{i,n} = \left(u_{i,n}(x); m \cos \frac{2\rho_{i,n}}{3} \cos \frac{\rho_{i,n}}{3} \right), \quad i = 1, 2; n \in Z^+.$$

Proof. To prove the first part of the lemma we take $(u, \alpha) \in L_p(0, 1) \oplus C$ and we define the functional $F(\widehat{u})$ as follows:

$$F(\widehat{u}) = mu \left(\frac{1}{3} \right) - \alpha.$$

Let us assume

$$U_\nu(\widehat{u}) = U_\nu(u), \nu = 1, 2, 3.$$

Then $F, U_\nu, \nu = 1, 2, 3$ are bounded linear functionals on $W_p^2(0, \frac{1}{3}) \cup (\frac{1}{3}, 1) \oplus C$, but unbounded on $L_p(0, 1) \oplus C$. Therefore (see, for example, [12, pp. 27-29]) the set

$$D(\mathcal{L}) = \left\{ \widehat{u} = (u, \alpha), u \in W_p^2 \left(0, \frac{1}{3} \right) \cup \left(\frac{1}{3}, 1 \right), F(\widehat{u}) = U_\nu(\widehat{u}) = 0, \nu = 1, 2, 3 \right\}.$$

is everywhere dense in $L_p(0, 1) \oplus C$, and \mathcal{L} is a closed operator as contraction of corresponding closed maximal operator.

The second part of the lemma is verified directly.

The lemma is proved.

For construction resolvent of operator \mathcal{L} , consider the equation

$$\mathcal{L} \widehat{u} - \lambda \widehat{u} = \widehat{f}, \quad (22)$$

where $\widehat{u} \in D(\mathcal{L})$, $\widehat{f} = (f, \beta) \in L_p(0, 1) \oplus C$. We can rewrite equation (22) in the form of

$$\begin{cases} -u'' = \lambda u + f, \\ u' \left(-\frac{1}{3} \right) - u' \left(\frac{1}{3} \right) - \lambda mu \left(\frac{1}{3} \right) = \beta, \\ U_\nu(u) = 0, \nu = 1, 2, 3. \end{cases} \quad (23)$$

Lemma 4. *For solution $\widehat{u} = (u, mu \left(\frac{1}{3} \right))$ of the equation (22) it holds the following representations*

$$u(x, \rho) = \frac{\beta \rho^2}{\Delta(\rho)} \left(e^{\frac{2i\rho}{3}} + e^{-\frac{2i\rho}{3}} \right) (e^{i\rho x} + e^{-i\rho x}) +$$

$$+ \int_0^{\frac{1}{3}} G_{11}(x, \rho, \xi) f(\xi) d\xi + \int_{\frac{1}{3}}^1 G_{12}(x, \xi, \rho) f(\xi) d\xi \quad (24)$$

if $x \in [0, \frac{1}{3}]$.

$$u(x, \rho) = \frac{\beta \rho^2}{\Delta(\rho)} \left(e^{\frac{i\rho}{3}} + e^{-\frac{i\rho}{3}} \right) \left(e^{i\rho(x-1)} + e^{-i\rho(x-1)} \right) + \int_0^{\frac{1}{3}} G_{21}(x, \rho, \xi) f(\xi) d\xi + \int_{\frac{1}{3}}^1 G_{22}(x, \xi, \rho) f(\xi) d\xi \quad (25)$$

if $x \in [\frac{1}{3}, 1]$

$$u\left(\frac{1}{3}, \rho\right) = \frac{\beta \rho^2}{\Delta(\rho)} \left(e^{\frac{2i\rho}{3}} + e^{-\frac{2i\rho}{3}} \right) \left(e^{\frac{i\rho}{3}} + e^{-\frac{i\rho}{3}} \right) + \int_0^{\frac{1}{3}} G_{21}(x, \rho, \xi) f(\xi) d\xi + \int_{\frac{1}{3}}^1 G_{22}(x, \rho, \xi) f(\xi) d\xi. \quad (26)$$

Proof. The solution of (23) will be sought in the form

$$u(x, \rho) = \begin{cases} C_{11}y_{11}(x) + C_{12}y_{12}(x) + y_1(x), & \text{for } x \in [0, \frac{1}{3}], \\ C_{21}y_{21}(x) + C_{22}y_{22}(x) + y_2(x), & \text{for } x \in [\frac{1}{3}, 1], \end{cases} \quad (27)$$

where $y_1(x)$ and $y_2(x)$ are defined by (14). Since $y(x)$, defined by (11) satisfies boundary conditions (2), then

$$U_\nu(y) = 0, \nu = \overline{1, 4}. \quad (28)$$

Let's demand the function $u(x, \rho)$ satisfy boundary conditions $U_\nu(u) = 0, \nu = 1, 2, 3, U_\nu(u) = \beta$. Then taking into account (28) from (27) we obtain

$$\begin{cases} C_{11}U_{\nu 1}(y_{11}) + C_{12}U_{\nu 1}(y_{12}) + C_{21}U_{\nu 2}(y_{21}) + C_{22}U_{\nu 2}(y_{22}) = 0, \nu = 1, 2, 3; \\ C_{11}U_{41}(y_{11}) + C_{12}U_{41}(y_{12}) + C_{21}U_{42}(y_{21}) + C_{22}U_{42}(y_{22}) = \beta. \end{cases}$$

Solving this system with respect to unknowns C_{kj} , we get

$$C_{11} = -\frac{\beta}{\Delta(\rho)} U_{11}(y_{12}) [U_{22}(y_{21})U_{32}(y_{22}) - U_{22}(y_{22})U_{32}(y_{21})],$$

$$C_{12} = \frac{\beta}{\Delta(\rho)} U_{11}(y_{11}) [U_{22}(y_{21})U_{32}(y_{22}) - U_{22}(y_{22})U_{32}(y_{21})],$$

$$C_{21} = \frac{\beta}{\Delta(\rho)} U_{22}(y_{22}) [U_{11}(y_{11})U_{31}(y_{12}) - U_{11}(y_{12})U_{31}(y_{11})],$$

$$C_{22} = -\frac{\beta}{\Delta(\rho)} U_{22}(y_{21}) [U_{11}(y_{11})U_{31}(y_{12}) - U_{11}(y_{12})U_{31}(y_{11})].$$

Substituting the received values of the coefficients C_{kj} in (27) and taking into account formula (8), (16)-(19), we obtain the validity of formulas (24) and (25). And formula (26) is obtained from (25) (or from (24)) by substitution $x = \frac{1}{3}$.

Lemma is proved.

4. Completeness of the eigenfunctions in spaces $L_p(0, 1) \oplus C$ and $L_p(0, 1)$

Theorem 2. *System $\{\hat{u}_i, n\}_{i=1,2;n \in Z^+}$ of eigenvectors of the operator \mathcal{L} is complete in $L_p(0, 1) \oplus C$, $1 < p < \infty$.*

Proof. To prove the completeness of the system of eigenfunctions of the operator \mathcal{L} in $L_p(0, 1) \oplus C$ we need to get the estimation of the resolvent of the operator \mathcal{L} at great values of $|\rho|$. We will use the following known inequalities

$$|\sin \rho| \leq ce^{|\rho| \sin \varphi}, |\cos \rho| \leq ce^{|\rho| \sin \varphi}, \tag{29}$$

where $\rho = re^{i\varphi}$, $0 \leq \varphi \leq \pi$. Besides, outside of circles of the same radius δ with centres in zero of $\sin \rho$ the following estimation is true

$$|\sin \rho| \geq m_\delta e^{r \sin \varphi}. \tag{30}$$

From estimations (29), (30) and from the formula (9) it follows that at great values of $|\rho|$ outside circles $K_{j,n}(\delta) = \{\rho: |\rho - \rho_{j,n}| < \delta\}$ of radius δ with the centres in zero of $\Delta(\rho)$ the following estimation is true

$$|\Delta(\rho)| \geq M_\delta r e^{r \sin \varphi}. \tag{31}$$

Assume $G(\delta) = C \setminus \bigcup_{j,n} K_{j,n}(\delta)$. From the representations (24), (25) considering the inequalities (29) – (31) we obtain the inequality

$$|u(x, \rho)| \leq \frac{C_\delta}{|\rho|}, \rho \in G(\delta), |\rho| \geq r_0,$$

which is fairly uniform on $x \in [0, 1]$. From the last estimation it follows that for the resolvent $R(\lambda) = (\mathcal{L} - \lambda I)^{-1}$ of the operator \mathcal{L} outside of the above-stated circles the following estimation is true

$$\|R(\rho^2)\| \leq \frac{C_\delta}{|\rho|}, |\rho| \geq r_0. \tag{32}$$

Having estimation (32), by a standard method (see for example [16]) we obtain that eigenfunctions of operator \mathcal{L} form a complete system in $L_p(0, 1) \oplus C$.

Let us note that the system $\{\hat{u}_{i,n}\}_{i=1,2;n \in Z^+}$ of eigenvectors of the operator \mathcal{L} has a biorthogonal-conjugate system $\{\hat{v}_{i,n}\}_{i=1,2;n \in Z^+}$, that are the system of eigenvectors of the conjugate operator \mathcal{L}^* , which in its turn is the linearized operator of the conjugate spectral problem:

$$v'' + \lambda v = 0, x \in \left(0, \frac{1}{3}\right) \cup \left(\frac{1}{3}, 1\right), (1^*)$$

$$v'(0) = v'(1) = 0,$$

$$v\left(\frac{1}{3} - 0\right) = v\left(\frac{1}{3} + 0\right), (2^*)$$

$$v' \left(\frac{1}{3} - 0 \right) - v' \left(\frac{1}{3} + 0 \right) = \lambda \bar{m} v \left(\frac{1}{3} \right).$$

Taking into account this by Theorem 2 we obtain

Corollary. System $\{\hat{u}_i, n\}_{i=1,2;n \in Z^+}$ of the eigenvectors of the operator \mathcal{L} is complete and minimal in $L_p(0, 1) \oplus C$, $1 < p < \infty$.

Now let us consider the completeness and minimality of a system $\{u_i, n\}_{i=1,2;n \in Z^+}$ of eigenfunctions of the problem (1), (2). It is clear that this system is overflowing in $L_p(0, 1)$: One function of this system is unnecessary. Let us clarify the following question: Which function can be excluded from the system while maintaining the properties of completeness and minimality and which can not? The answer to this question is given by the following theorem.

Theorem 3. Let n_0 be some number from the set of indexes Z^+ . Then the system obtained from the system $\{\hat{u}_i, n\}_{i=1,2;n \in Z^+}$ by excluding on arbitrary function u_{i,n_0} is complete and minimal in the space $L_p(0, 1)$, $1 < p < \infty$

Proof. According to the above mentioned, the system $\{u_i, n\}_{i=1,2;n \in Z^+}$ has a biorthogonal-conjugated system $\{\hat{v}_i, n\}_{i=1,2;n \in Z^+}$, where

$$\hat{v}_n = \left(v_n(x); \bar{m} v_n \left(\frac{1}{3} \right) \right),$$

and $v_n(x)$ are eigenfunctions of the adjoint problem (1*), (2*). Carrying out similar calculations for eigenfunctions, we find that the formulas are valid

$$v_{i,n}(x) = \begin{cases} c_{i,n} \cos \frac{2\rho_{i,n}}{3} \cos \rho_{i,n} x, & x \in \left[0, \frac{1}{3}\right], \\ c_{i,n} \cos \frac{\rho_{i,n}}{3} \cos \rho_{i,n} (1-x), & x \in \left[\frac{1}{3}, 1\right], \end{cases} \quad i = 1, 2; n \in Z^+$$

where $c_{i,n}$ – are normalization numbers. From this formula it is clear that if n_0 is number, from the set of index Z^+ then $v_{i,n_0} \left(\frac{1}{3} \right) \neq 0$. Therefore, all statements of the theorem follow from the results of [17] (see also [18]).

References

- [1] A.N. Tikhonov, A.A. Samarskii, *Equations of mathematical physics*, Courier Corporation, 2013.
- [2] F.V. Atkinson, *Discrete and continuous boundary problems*, Academic Press, New York, London, 1964.
- [3] L. Collatz, *Eigenwertaufgaben mit technischen Anwendungen*, Leipzig, 1963
- [4] T.B. Gasymov, Sh.J. Mammadova, *On convergence of spectral expansions for one discontinuous problem with a spectral parameter in the boundary condition*, Trans.of NAS of Azerb., **XXVI**(4), 103–116.

- [5] T.B. Gasymov, A.A. Huseynli, *On completeness of eigenfunctions and associated functions of a discontinuous second order differential operator*, Dokl. NAN of Azerb., **LXVIII**(1), 3–7.
- [6] N.Yu. Kapustin, E.I. Moiseev, *The basis property in L_p of the systems of eigenfunctions corresponding to two problems with a spectral parameter in the boundary condition*, Differential equations, **36**(10), 2000, 1498–1501.
- [7] N.Yu. Kapustin, E.I. Moiseev, *On spectral problems with a spectral parameter in the boundary condition*, Diff. Uravn., **33**(1), 1997, 16–21.
- [8] N.B. Kerimov, V.S. Mirzoyev, *On basis properties of one spectral problem with spectral parameter in the boundary condition*, Sib. Math. Jour., 2003,**44**(5), 1041–1045. (Russian)
- [9] N.B. Kerimov, R.G. Poladov, *On basicity in $L_p(0,1)$, ($1 < p < \infty$) of the system of eigenfunctions of one boundary value problem. II*. Proc. of IMM of NAS of Azerb., **XXIII**, 2005, 65–76.
- [10] B.M. Levitan, I.S. Sargsjan, *Introduction to spectral theory*, American Mathematical Society, 1975.
- [11] M.A. Naimark, *Linear Differential operators*, Frederik Ungar Publ. Co., New York, 1968.
- [12] V.E. Lyantse, O.G. Storozh, *Methods of the theory of unbounded operators*, Kiev, 1983.
- [13] N. Dunford, J.T. Schwartz, *Linear operators*, Part 3, New York, London, Sydney, Toronto, 1971.
- [14] T.B. Gasymov, *On convergence of spectral expansions for one discontinuous problem with a spectral parameter in the boundary condition*, Trans. of NAS of Azerb., **XXVI**(4), 103–116.
- [15] T.B. Gasymov, T.Z. Garayev, *On convergence of spectral expansions for one discontinuous problem with spectral parameter in the boundary condition*, Trans. of NAS of Azerb., **XXVI**(4), 103–116.
- [16] B.T. Bilalov, T.R. Muradov, *Defective bases of Banach spaces*, Proc. IMM NAS Azerb., **XXII**, 2005, 23–26.

Alirza Q. Akhmedov
Baku State University, Az1148, Baku, Azerbaijan
E-mail: ehmedov-elirza@mail.ru

Ilkin Q. Feyzullayev

Institute of Mathematics and Mechanics of NAS of Azerbaijan, Az1141, Baku, Azerbaijan

E-mail: ilkin0704@mail.ru

Received 05 September 2022

Accepted 22 October 2022