Caspian Journal of Applied Mathematics, Ecology and Economics V. 12, No 2, 2023, December ISSN 1560-4055

Nodal Solutions of Some Nonlinear Dirac Problems

Humay Sh. Rzayeva

Abstract. In this paper we consider the nonlinear boundary value problem for the one-dimensional Dirac operator. We show the existence of pair of nodal solutions of this problem. The proof of our main result based on a method on bifurcation from zero and infinity.

Key Words and Phrases: nonlinear Dirac equation, eigenvalue, eigenvector-function, bifurcation from zero, bifurcation from infinity, nodal solution

2010 Mathematics Subject Classifications: 34A30, 34B15, 34C23, 34K29, 47J10, 47J15

1. Introduction

We consider the following nonlinear Dirac equation

$$(\ell w)(x) \equiv Bw'(x) - P(x)w(x) = sf(w(x)), \ 0 < x < \pi,$$
(1.1)

with the boundary conditions

$$U_1(w) := (\sin \alpha, \, \cos \alpha) \, w(0) = v(0) \cos \alpha + u(0) \sin \alpha = 0, \tag{1.2}$$

$$U_2(w) := (\sin\beta, \cos\beta)w(\pi) = v(\pi)\cos\beta + u(\pi)\sin\beta = 0, \qquad (1.3)$$

where

$$B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, P(x) = \begin{pmatrix} p(x) & 0 \\ 0 & r(x) \end{pmatrix}, w(x) = \begin{pmatrix} u(x) \\ v(x) \end{pmatrix},$$

 $\lambda \in \mathbb{C}$ is an eigenvalue parameter, $p(x), r(x) \in C([0,\pi];\mathbb{R}), \alpha, \beta$, are real constants such that $0 \leq \alpha, \beta < \pi$. Here s is a nonzero real number and the function $f = \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} \in C(\mathbb{R}^2 \times \mathbb{R}^2; \mathbb{R}^2)$ satisfies the following condition:

$$f(w) = \gamma w + h(w) \text{ and } f(w) = \delta w + \hbar(w), \qquad (1.4)$$

where $\gamma, \delta \ (\gamma \neq \delta)$ are positive constants and

http://www.cjamee.org

(C) 2013 CJAMEE All rights reserved.

Humay Sh. Rzayeva

$$h(w) = \begin{pmatrix} h_1(w) \\ h_2(w) \end{pmatrix} = o(|w|) \text{ as } |w| \to 0,$$
(1.5)

$$\hbar(w) = \begin{pmatrix} \hbar_1(w) \\ \hbar_2(w) \end{pmatrix} = o(|w|) \text{ as } |w| \to \infty$$
(1.6)

(here by $|\cdot|$ we denote the norm in \mathbb{R}^2).

The Dirac equations describing spin-1/2 particles, such as electrons and positrons, play the important role in both physics and mathematics. The nonlinear Dirac equations describes the self-action in quantum electrodynamics and used as an effective theory in atomic and gravitational physics (see [14]).

The existence of nodal solutions of boundary value problems for ordinary differential operators of second and fourth orders was study in [5-11]. In these papers, the authors, using the bifurcation technique, show the existence of solutions of problems they consider, contained in classes of functions with a fixed usual oscillation count.

In this paper, we will show the existence of nodal solutions of problem (1.1)-(1.3) using global bifurcation results from zero and infinity that we previously obtained for nonlinear Dirac eigenvalue problems.

2. Preliminary

Let (b.c.) be denote the set of vector-functions that satisfies the boundary conditions (1.2) and (1.3).

We consider the following nonlinear eigenvalue problem

$$\begin{cases} (\ell w)(x) = \lambda w(x) + g(x, w(x), \lambda), \ x \in (0, \pi), \\ w \in (b.c.). \end{cases}$$

$$(2.1)$$

We will impose one or the other of the following conditions on the nonlinear term $g(x, w, \lambda) \in C([0, \pi] \times \mathbb{R}^2 \times \mathbb{R}; \mathbb{R}^2)$:

$$g(x, w, \lambda) = o(|w|) \text{ as } |w| \to 0, \qquad (2.2)$$

or

$$g(x, w, \lambda) = o(|w|) \text{ as } |w| \to +\infty,$$
(2.3)

uniformly with respect to $(x, \lambda) \in [0, \pi] \times \Lambda$, for any bounded interval $\Lambda \subset \mathbb{R}$.

By conditions (2.2) and (2.3) the nonlinear problem (2.1) is linearizable both at zero and at infinity, and the corresponding linear problem is

$$\begin{cases} (\ell w)(x) = \lambda w(x), \ x \in (0, \pi), \\ w \in (b.c.). \end{cases}$$
(2.4)

Let *E* be the Banach space $C([0, \pi]; \mathbb{R}^2) \cap \{w : U(w) = 0\}$ with the norm $||w||_0 = \max_{x \in [0, \pi]} |u(x)| + \max_{x \in [0, \pi]} |v(x)|$. We denote by *S* a subset of *E*, which defined as follows:

$$S = \{ w \in E : |u(x) + |v(x)| > 0, \, \forall x \in [0, \pi] \}.$$

As in [2] (see also [4]), for each $w = \begin{pmatrix} u \\ v \end{pmatrix} \in S$ we define a continuous function $\theta(w, x)$ on $[0, \pi]$ by

$$\tan \theta(w, x) = \frac{v(x)}{u(x)}, \ \theta(w, 0) = -\alpha.$$
(2.5)

Theorem 2.1 [2, Theorem 3.1]. The eigenvalues $\lambda_k, k \in \mathbb{Z}$, of the problem (2.4) are real and simple, and can be numbered in ascending order on the real axis

$$\ldots \ <\lambda_{-k}< \ \ldots \ <\lambda_{-1}<\lambda_0<\lambda_1< \ \ldots \ <\lambda_k< \ \ldots \ ,$$

so that for the eigenvector-function $w_k(x)$, $k \in \mathbb{Z}$, corresponding to the eigenvalue λ_k , the angular function $\theta(w_k, x)$ at $x = \pi$ satisfy the condition

$$\theta(w_k, \pi) = -\beta + k\pi. \tag{2.6}$$

The eigenvector-functions $w_k(x) = \begin{pmatrix} u_k(x) \\ v_k(x) \end{pmatrix}$ have, with a suitable interpretation, the following oscillation properties: if k > 0 and $k = 0, \alpha \ge \beta$ (except the cases $\alpha = \beta = 0$ and $\alpha = \beta = \pi/2$), then

$$\begin{pmatrix} s(u_k) \\ s(v_k) \end{pmatrix} = \begin{pmatrix} k - 1 + \chi(\alpha - \pi/2) + \chi(\pi/2 - \beta) \\ k - 1 + \operatorname{sgn}\alpha \end{pmatrix},$$
(2.7)

and if k < 0 and k = 0, $\alpha < \beta$, then

$$\begin{pmatrix} s(u_k) \\ s(v_k) \end{pmatrix} = \begin{pmatrix} |k| - 1 + \chi(\pi/2 - \alpha) + \chi(\beta - \pi/2) \\ |k| - 1 + \operatorname{sgn}\beta \end{pmatrix},$$
(2.8)

where s(g) the number of zeros of the function $g \in C([0,\pi]; \mathbb{R})$ in the interval $(0,\pi)$ and

$$\chi(x) = \begin{cases} 0, & \text{if } x \le 0, \\ 1, & \text{if } x > 0. \end{cases}$$

Moreover, the functions $u_k(x)$ and $v_k(x)$ have only nodal zeros in the interval $(0,\pi)$.

For each $k \in \mathbb{Z}$ and each ν , let S_k^{ν} denote the set of vector functions $w \in S$ with the following properties:

(i) $\theta(w,\pi) = -\beta + k\pi;$

(ii) if k > 0 or $k = 0, \alpha \ge \beta$ (except the cases $\alpha = \beta = 0$ and $\alpha = \beta = \pi/2$), then for fixed w, as x increases from 0 to π , the function θ cannot tend to a multiple of $\pi/2$ from above, and as x decreases, the function θ cannot tend to a multiple of $\pi/2$ from below; if

k < 0 or $k = 0, \alpha < \beta$, then for fixed w, as x increases, the function θ cannot tend to a multiple of $\pi/2$ from below, and as x decreases, the function θ cannot tend to a multiple of $\pi/2$ from above;

(iii) the function $\nu u(x)$ is positive in a deleted neighborhood of x = 0.

Remark 2.1. By (2.5), Theorem 2.1 and [2, Remark 2.1] we have $w_k \in S_k = S_k^- \cup S_k^+$, in other words the sets S_k^- , S_k^+ and S_k are nonempty. From the definition of the sets S_k^- , S_k^+ and S_k it can be seen that they are disjoint and open in E.

Now for problem (1.1)-(1.3) we can give global bifurcation results obtained earlier in [1, 3].

Theorem 2.2 [3, Theorem 3.1]. Let condition (2.2) holds. Then for each $k \in \mathbb{Z}$ and each ν , there exists a continuum C_k^{ν} of solutions of problem (2.1) which contains $(\lambda_k, \tilde{0})$, is lies in $(\mathbb{R} \times S_k^{\nu}) \cup \{(\lambda_k, \tilde{0})\}$ and is unbounded in $\mathbb{R} \times E$.

Theorem 2.3 [1, Theorem 1]. Let condition (2.3) holds. Then for each $k \in \mathbb{Z}$ and each ν there exists a continua D_k^{ν} of solutions of problem (2.1) which contains (λ_k, ∞) and has the following properties:

(i) there exists a neighborhood V_k of (λ_k, ∞) in $\mathbb{R} \times E$ such that $V_k \cap D_k^{\nu} \subset \mathbb{R} \times S_k^{\nu}$;

(ii) either D_k^{ν} meets (λ'_k, ∞) with respect to the set $\mathbb{R} \times S_{k'}^{\nu'}$ for some $(\tilde{k'}, \nu') \neq (\tilde{k}, \nu)$, or D_k^{ν} meets $(\lambda, \tilde{0})$ for some $\lambda \in \mathbb{R}$, or D_k^{ν} has an unbounded projection onto $\mathbb{R} \times \{\tilde{0}\}$.

3. Existence of nodal solutions to the nonlinear problem (1.1)-(1.3)

In this section, for each $k \in \mathbb{Z}$ and each ν , an interval is determined for s in which there are solutions to the problem (1.1)-(1.3) contained in the set S_k^{ν} .

Theorem 1. Let $\lambda_k \neq 0$ and

$$\frac{\lambda_k}{\gamma} < s < \frac{\lambda_k}{\delta} \ \text{ or } \ \frac{\lambda_k}{\delta} < s < \frac{\lambda_k}{\gamma}$$

for some $k \in \mathbb{Z}$. Then problem has two solutions $w_{k,+}$ and $w_{k,-}$ such that $w_{k,+} \in S_k^+$ and $w_{k,-} \in S_k^-$.

Proof. By the first relation of (1.5) we can rewrite (1.1)-(1.3) it following form

$$\begin{cases} (\ell w)(x) = s\gamma w(x) + sh(w(x)), \ x \in (0,\pi), \\ w \in (b.c.). \end{cases}$$
(3.2)

Alongside problem (3.2) we shall consider the following nonlinear eigenvalue problem

$$\begin{cases} (\ell w)(x) = \lambda s \gamma w(x) + s h(w(x)), \ x \in (0, \pi), \\ w \in (b.c.), \end{cases}$$
(3.3)

Nodal Solutions of Some Nonlinear Dirac Problems

or, equivalently

$$\begin{cases} (\tilde{\ell}w)(x) = \lambda w(x) + \tilde{g}(x,w), \ x \in (0,\pi), \\ w \in (b.c.), \end{cases}$$
(3.4)

where

$$\tilde{\ell}w = \frac{1}{s\gamma}\ell w$$
 and $\tilde{g}(x,w) = \frac{1}{\gamma}h(w(x)).$

In view of condition (1.5) we have

$$\tilde{g}(x,w) = o(|w|) \text{ as } |w| \to 0,$$
(3.5)

uniformly with respect to $x \in [0, \pi]$. Then, in view of (3.5), by Theorem 2.2 for each $k \in \mathbb{Z}$ and each ν , there exists a continuum \tilde{C}_k^{ν} of solutions of problem (3.4) (also problem (3.3)) which contains $(\tilde{\lambda}_k, \tilde{0})$, is lies in $(\mathbb{R} \times S_k^{\nu}) \cup \{(\tilde{\lambda}_k, \tilde{0})\}$ and is unbounded in $\mathbb{R} \times E$, where $\tilde{\lambda}_k$ is a *k*th eigenvalue of the linear problem

$$\begin{cases} (\tilde{\ell}w)(x) = \lambda w(x), \ x \in (0,\pi), \\ w \in (b.c.), \end{cases}$$

also of the linear problem

$$\begin{cases} (\ell w)(x) = \lambda s \gamma w(x), \ x \in (0, \pi), \\ w \in (b.c.). \end{cases}$$
(3.6)

In view of (3.6) we get

$$\tilde{\lambda}_k = \frac{\lambda_k}{s\gamma}.$$

By the second representation of (1.4) problem (3.3) will take the form

$$\begin{cases} (\ell w)(x) - s(\delta - \gamma)w(x) = \lambda s \gamma w(x) + s \hbar(w(x)), \ x \in (0, \pi), \\ w \in (b.c.), \end{cases}$$
(3.6)

or, equivalently

$$\begin{cases} (\hat{\ell}w)(x) = \lambda w(x) + \hat{g}(x, w), \ x \in (0, \pi), \\ w \in (b.c.), \end{cases}$$
(3.7)

where

$$\hat{\ell}w = \frac{1}{s\gamma}\ell w - \left(\frac{\delta}{\gamma} - 1\right)w$$
 and $\hat{g}(x, w) = \frac{1}{\gamma}h(w(x)).$

In view of condition (1.6) for any bounded intervals $\Lambda \subset \mathbb{R}$

$$\hat{g}(x, w, \lambda) = o(|w|) \text{ as } |w| \to +\infty,$$
(3.8)

uniformly with respect to $(x, \lambda) \in [0, \pi] \times \Lambda$. Then, by (3.8), it follows from Theorem 2.3 that for each $k \in \mathbb{Z}$ and each ν there exists a continua \hat{C}_k^{ν} of solutions of problem (3.7) (also of problems (3.7) and (3.2)) which contains $(\hat{\lambda}_k, \infty)$ and has the following properties:

37

Humay Sh. Rzayeva

(i) there exists a neighborhood \hat{V}_k of $(\hat{\lambda}_k, \infty)$ in $\mathbb{R} \times E$ such that $\hat{V}_k \cap \hat{C}_k^{\nu} \subset \mathbb{R} \times S_k^{\nu}$;

(ii) either \hat{C}_k^{ν} meets $(\hat{\lambda}'_k, \infty)$ with respect to the set $\mathbb{R} \times S_{k'}^{\nu'}$ for some $(k', \nu') \neq (k, \nu)$, or \hat{C}_k^{ν} meets $(\lambda, \tilde{0})$ for some $\lambda \in \mathbb{R}$, or \hat{C}_k^{ν} has an unbounded projection onto $\mathbb{R} \times \{\tilde{0}\}$, where $\hat{\lambda}_k$ is a *k*th eigenvalue of the linear problem

$$\begin{cases} (\hat{\ell}w)(x) = \lambda w(x), \ x \in (0,\pi), \\ w \in (b.c.), \end{cases}$$

also of the linear problem

$$\begin{cases} (\ell w)(x) = s(\lambda \gamma + \delta - \gamma)w(x) + s\hbar(w(x)), \ x \in (0, \pi), \\ w \in (b.c.). \end{cases}$$
(3.9)

By (3.9) we get

$$\hat{\lambda}_k = \frac{\lambda_k}{s\gamma} - \left(\frac{\delta}{\gamma} - 1\right).$$

Since conditions (1.5) and (1.6) are satisfied simultaneously, it follows from the proof of the first part of [13, Theorem 3.3] that $\hat{C}_k^{\nu} \subset \mathbb{R} \times S_k^{\nu}$, and consequently, the first alternative of property (ii) of \hat{C}_k^{ν} cannot hold, i.e., \hat{C}_k^{ν} cannot meets (λ'_k, ∞) for any $k' \neq k$. Then either \hat{C}_k^{ν} meets $(\lambda, 0)$ for some $\lambda \in \mathbb{R}$, or the projection \hat{C}_k^{ν} onto $\mathbb{R} \times \{\tilde{0}\}$ is unbounded.

Let \hat{C}_k^{ν} meets $(\lambda, \tilde{0})$ for some $\lambda \in \mathbb{R}$. Then, using [12, Lemma 1.24], we can show that $\lambda = \tilde{\lambda}_k$. Moreover, if \tilde{C}_k^{ν} meets (λ, ∞) for some $\lambda \in \mathbb{R}$, then from property (i) of \hat{C}_k^{ν} it follows that $\lambda = \hat{\lambda}_k$.

If the projection \hat{C}_k^{ν} onto $\mathbb{R} \times \{\tilde{0}\}$ is unbounded, then there exists $\{(\mu_n, z_n)\}_{n=1}^{\infty} \subset (\hat{C}_k^{\nu} \setminus \{(\hat{\lambda}_k, \infty)\}) \subset \mathbb{R} \times S_k^{\nu} \left(z_n = \begin{pmatrix} z_{1n} \\ z_{2n} \end{pmatrix}\right)$ such that either $\lim_{n \to \infty} \mu_n \to +\infty \text{ or } \lim_{n \to \infty} \mu_n \to -\infty.$

We introduce the notations:

$$\varphi_{n,1}(x) = \frac{\hbar_1(z_n(x))z_{n,1}(x)}{z_{n,1}^2(x) + z_{n,2}^2(x)}, \quad \psi_{n,1}(x) = \frac{\hbar_1(z_n(x))z_{n,2}(x)}{z_{n,2}^2(x) + z_{n,2}^2(x)}$$

$$\varphi_{n,2}(x) = \frac{\hbar_2(z_n(x))z_{n,1}(x)}{z_{n,1}^2(x) + z_{n,2}^2(x)}, \quad \psi_{n,2}(x) = \frac{\hbar_2(z_n(x))z_{n,1}(x)}{z_{n,1}^2(x) + z_{n,2}^2(x)}.$$
(3.10)

Let $\epsilon_0 > 0$ be a fixed small number. Then, by conditions (1.5) and (1.6), there exist a small $\rho_0 > 0$ and a large $\rho_0 > 0$ such that

$$\frac{|h(w)|}{|w|} < \epsilon_0 \text{ for any } w \in S, \ |w| < \rho_0, \tag{3.11}$$

$$\frac{|\hbar(w)|}{|w|} < \epsilon_0 \text{ for any } w \in S, \ |w| > \varrho_0.$$
(3.12)

38

Since $\hbar \in C(\mathbb{R}^2 \times \mathbb{R}^2; \mathbb{R}^2)$ there exists $K_0 > 0$ such that

$$\frac{|\hbar(w)|}{|w|} < K_0 \text{ for any } w \in E, \ \rho_0 \le |w| \le \varrho_0.$$
(3.13)

By (1.4) and (3.11) for any $w \in S$ such that $|w| < \rho_0$ we get

$$|\hbar(w)| = |(\gamma - \delta)w + h(w)| \le (|\gamma - \delta| + \epsilon_0)|w|.$$

$$(3.14)$$

Thus, it follows from (3.12)-(3.14) that

$$|\hbar(w)| \le M|w|, \ w \in S,\tag{3.15}$$

where

$$M = \max \{ K_0, |\gamma - \delta| + \epsilon_0 \}.$$

In view of (3.15), by (3.10) we get

$$\begin{aligned} |\varphi_{n,1}(x)| &\leq \frac{|\hbar_1(z_n(x))| \, |z_{n,1}(x)|}{z_{n,1}^2(x) + z_{n,2}^2(x)} \leq 2M, \quad |\psi_{n,1}(x)| \leq \frac{|\hbar_1(z_n(x))| \, |z_{n,2}(x)|}{z_{n,1}^2(x) + z_{n,2}^2(x)} \leq 2M, \\ |\varphi_{n,2}(x)| &\leq \frac{|\hbar_2(z_n(x))| \, |z_{n,1}(x)|}{z_{n,1}^2(x) + z_{n,2}^2(x)} \leq 2M, \quad |\psi_{n,2}(x)| \leq \frac{|\hbar_2(z_n(x))| \, |z_{n,2}(x)|}{z_{n,1}^2(x) + z_{n,2}^2(x)} \leq 2M. \end{aligned}$$
(3.16)

By (3.10) it follows from (3.9) that $(\mu_n, z_n), n \in \mathbb{N}$, solves the linear problem

$$\begin{cases} (\ell w)(x) - sP_n(x)w(x) = s(\lambda\gamma + \delta - \gamma)w(x), \ x \in (0,\pi), \\ w \in (b.c.), \end{cases}$$
(3.17)

where

$$P_n(x) = \left(\begin{array}{cc} \varphi_{n,1}(x) & \psi_{n,1}(x) \\ \varphi_{n,2}(x) & \psi_{n,2}(x) \end{array}\right).$$

By [3, theorem 2.4] we have

 $\theta'(z_n, x) =$

$$s(\mu_n \gamma + \delta - \gamma) + p(x) \cos^2 \theta(z_n, x) + r(x) \sin^2 \theta(z_n, x) + s\varphi_{n,1}(x) \cos^2 \theta(z_n, x) + s\psi_{n,2}(x) \sin^2 \theta(z_n, x) + \frac{1}{2}s \{\psi_{n,1}(x) + \varphi_{n,2}(x)\} \sin 2\theta(z_n, x).$$
(3.18)

Integrating (3.18) from 0 to π and using (2.5), (2.6) we get

$$s(\mu_{n}\gamma + \delta - \gamma) = k\pi + \alpha - \beta - \int_{0}^{\pi} \left\{ p(x)\cos^{2}\theta(z_{n}, x) + r(x)\sin^{2}\theta(z_{n}, x) \right\} dx - \left\{ \int_{0}^{\pi} \left\{ \delta + \varphi_{n,1}(x)\cos^{2}\theta(z_{n}, x) + \psi_{n,2}(x)\sin^{2}\theta(z_{n}, x) \right\} dx + \frac{1}{2} \int_{0}^{\pi} \left\{ \psi_{n,1}(x) + \varphi_{n,2}(x) \right\} \sin 2\theta(z_{n}, x) dx \right\}.$$
(3.19)

39

Humay Sh. Rzayeva

The left side of (3.19) for sufficiently large n takes on sufficiently large values in absolute value, while by $p, r \in C([0, \pi]; \mathbb{R})$ and (3.16) the right side remains bounded, which gives a contradiction. Therefore, the projection of \hat{C}_k^{ν} onto $\mathbb{R} \times \{\tilde{0}\}$ is bounded, and consequently, \hat{C}_k^{ν} meets $(\tilde{\lambda}_k, 0)$. Similarly, we can show that the projection of \tilde{C}_k^{ν} onto $\mathbb{R} \times \{\tilde{0}\}$ is also bounded, which implies that \tilde{C}_k^{ν} meets $(\hat{\lambda}_k, \infty)$. Therefore, for $k \in \mathbb{Z}$ and each ν we have

$$\tilde{C}_k^{\nu} = \hat{C}_k^{\nu}. \tag{3.20}$$

From (3.2) it is clear that the solution to this problem (λ, w) with $\lambda = 1$ is also solution to problem (1.1)-(1.3). Since \tilde{C}_k^{ν} is connected, it follows that for the existence of a solution $w \in E$ of problem (1.1)-(1.3) contained in S_k^{ν} for some $k \in \mathbb{Z}$, by (3.20), it is sufficient that on the real axis \mathbb{R} the point $\frac{\lambda_k}{s\gamma}$ lies to the left of 1 and the point $\frac{\lambda_k}{s\gamma} - (\frac{\delta}{\gamma} - 1)$ lies to the right of 1, or the point $\frac{\lambda_k}{s\gamma} - (\frac{\delta}{\gamma} - 1)$ lies to the right of 1, and the point $\frac{\lambda_k}{s\gamma}$ lies to the left of 1.

If $\lambda_k > 0$, s > 0 and $\frac{\lambda_k}{\gamma} < s < \frac{\lambda_k}{\delta}$, then we have

$$\frac{\lambda_k}{s\gamma} < 1 \text{ and } 1 < \frac{\lambda_k}{s\delta}.$$

By conditions s > 0, $\delta > 0$ and $\gamma > 0$ we get

$$1 < \frac{\lambda_k}{s\delta} \Longrightarrow s\delta < \lambda_k \Longrightarrow s\frac{\delta}{\gamma} < \frac{\lambda_k}{\gamma} \Longrightarrow s\left(1 + \frac{\delta}{\gamma} - 1\right) < \frac{\lambda_k}{\gamma} \Longrightarrow$$
$$s < \frac{\lambda_k}{\gamma} - s\left(\frac{\delta}{\gamma} - 1\right) \Longrightarrow 1 < \frac{\lambda_k}{s\gamma} - \left(\frac{\delta}{\gamma} - 1\right).$$

Therefore, we have the following relation

$$\frac{\lambda_k}{s\gamma} < 1 < \frac{\lambda_k}{s\gamma} - \left(\frac{\delta}{\gamma} - 1\right).$$

Other cases are considered similarly. The proof of this theorem is complete.

References

- Z.S. Aliyev ZS, N.A. Neymatov, H.Sh. Rzayeva, Unilateral global bifurcation from infinity in nonlinearizable one-dimensional Dirac problems. *Int. J. Bifur. Chaos.* 2021; 31(1): 1-10.
- [2] Z.S. Aliyev, H.Sh. Rzayeva, Oscillation properties for the equation of the relativistic quantum theory, Appl. Math. Comput. 271 (2015), 308-316.
- [3] Z.S. Aliyev, H.Sh. Rzayeva, Global bifurcation for nonlinear Dirac problems, *Electron. J. Qual. Theory Differ. Equ.* (46) (2016), 1-14.

- [4] F.V. Atkinson, Discrete and Continuous Boundary Problems, Academic Press, New York, London, 1964.
- [5] G. Dai, Global bifurcation from intervals for Sturm-Liouville problems which are not linearizable, *Elec. J. Qual. Theory of Diff. Equat.* (65) (2013), 1-7.
- [6] L.H. Erbe, H.Y. Wang, On the existence of positive solutions of ordinary differential equations, Proc. Amer. Math. Soc. 120(3) (1994), 743–748.
- [7] J. Henderson, H.Y. Wang, Positive solutions for nonlinear eigenvalu eproblems, J.Math.Anal.Appl. 208 (1997), 252-259.
- [8] R. Ma and G. Dai, Global bifurcation and nodal solutions for a Sturm-Liouville problem with a nonsmooth nonlinearity, J. Funct. Anal. 265 (2013), 1443-1459.
- [9] R. Ma, B.Thompson, Nodal solutions for nonlinear eigenvalue problems, Nonlinear Anal. 59 (2004), 707–718.
- [10] R. Ma, B. Thompson, Multiplicity results for second-order two-point boundary value problems with nonlinearities across several eigenvalues, *Appl. Math. Lett.* 18(5) (2005), 587-595.
- [11] Y. Naito, S. Tanaka, On the existence of multiple solutions of the boundary value problem for nonlinear second-order differential equations, *Nonlinear Anal.* 56(4) (2004), 919-935
- [12] P.H. Rabinowitz, Some global results for nonlinear eigenvalue problems, J. Funct. Anal. 7(3) (1971), 487-513.
- [13] B.P. Rynne, Bifurcation from zero or infinity in Sturm-Liouville problems which are not linearizable, J. Math. Anal. Appl. 228(1) (1998), 141-156.
- [14] B. Thaller, *The Dirac Equation*, Springer, Berlin, 1992.

Humay Sh. Rzayyeva Institute of Mathematics and Mechanics of Ministry of Science and Education of Azerbaijan, Baku, Azerbaijan E-mail: humay_rzayeva@bk.ru

Received 01 October 2023 Accepted 21 November 2023